

# Irrigation Water Quality

Lydia Zamlynny and Dr. Jennie Rand

## Background

Successful crop farming is often dependent on the quality of irrigation water that is readily available. Salinity, toxic ions, heavy metals and other parameters can affect the crops and soil health. Microbiological pathogens in the water can also present issues by contaminating crops and causing biofouling in irrigation equipment.

## Irrigation System Model




**Figure 1:** Constructed Irrigation System Model.

The purpose of this project is to evaluate microbial concentrations (*E. coli*) throughout an irrigation system considering biofilm and spray head pressure. An irrigation system was simulated using PVC pipe, an annular reactor (AR), hose, and sprinkler heads (shown in Figure 1). The AR was used to represent the pipeline through which water is delivered to an irrigation system, and the constructed pipes attempt to replicate the conditions expected in a 40-acre field system.

## Our Research

Feed water is pumped into the system at a specified rate, and *E. coli* K12 concentrations are monitored by plating samples taken at various stages in the system. Figure 2 shows the plates and growing colonies from a sample of the stock solution.



**Figure 2:** *E. coli* Stock Plates

The goal of the project will be to study the interactions between the system and the pathogens being introduced through the feed water. Two considerations under evaluation include:

- Formation of biofilm and subsequently microbiological regrowth on pipe surfaces, and how bacteria may re-enter the bulk water to be sprayed on crops.
- The potential for the pressure of the spray heads to damage microbiological cells as they exit the system.